WORLD INTELLECTUAL PROPERTY ORGANIZATION



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5: C07C 233/75, A61K 31/165 C07D 213/81, 213/82, C07C 233/29 C07C 235/56, C07D 213/75, 239/42

(11) International Publication Number:

WO 93/25517

(43) International Publication Date:

23 December 1993 (23.12.93)

(21) International Application Number:

PCT/GB93/01266

(22) International Filing Date:

15 June 1993 (15.06.93)

(74) Agent: SKAILES, Humphrey, J.; Frank B. Dehn & Co., Imperial House, 15-19 Kingsway, London WC2B 6UZ

(GB).

(30) Priority data:

9212673.9 9212693.7

15 June 1992 (15.06.92) GB 15 June 1992 (15.06.92) GB

(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).

(71) Applicant: CELLTECH LIMITED [GB/GB]; 216 Bath Road, Slough, Berkshire SL1 4EN (GB).

(72) Inventors: BEELEY, Nigel, Robert, Arnold; 16 Cheshire Road, Thame, Oxfordshire OX9 3LQ (GB). MILLI-CAN, Thomas, Andrew; 3 Harcourt Close, Dorney Reach, Maidenhead, Berkshire SL6 0DY (GB).

Published

With international search report.

(54) Title: TRISUBSTITUTED PHENYL DERIVATIVES AS SELECTIVE PHOSPHODIESTERASE IV INHIBITORS

$$X - R^2$$

$$Z(CH_2)_n R^4$$
(1)

(57) Abstract

Compounds of formula (i) are described wherein Y represents a halogen atom or a group -OR1 where R1 represents an optionally substituted alkyl group; R2 represents an optionally substituted cycloalkyl, cycloalkenyl or polycycloalkyl group; Z represents -N(R3)CO- or -CONE(R3) represents a hydrogen atom or an alkyl, aryl or aralkyl group; R4 represents an aryl or heteroaryl group; X represents -O-, -S-, -CH₂- or -N(\mathbb{R}^5)-, where \mathbb{R}^5 is a hydrogen atom or an alkyl group; n is zero or an integer or value 1, 2 or 3; and the salts, solvates and hydrates thereof. The compounds are selective phosphodiesterase IV inhibitors and are useful for the prophylaxis or treatment of inflammatory diseases.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

Austria	FR	France	MR	Mauritania
Australia	GA	Gabon	MW	Malawi
Barhados	GB	United Kingdom	NL	Netherlands
Belgium	GN	Guinea	NO	Norway
Burkina Faso	GR	Greece	NZ	New Zealand
Bulgaria	HU	Hungary	PL	Poland
Benin	31	Ireland	PT	Portugal
Brazil .	IT	łtoly	RO	Romania
Canada	JP	Japan	RU	Russian Federation
Central African Republic	KP	Democratic People's Republic	SD	Sudan
Congo		of Korea	SE	Sweden
Switzerland	KR	Republic of Korea	SK	Slovak Republic
Côte d'Ivoire	KZ	Kazakhstan	SN	Senegal
Cameroon	LI	Licehtenstein	SU	Soviet Union
Czechoslovakia ·	LK	Sri Lanka .	TD	Chad
Czech Republic	1.0	Luxembourg	TG	T'ogo
Germany	MC	Monaco	UA	Ukraine
Denmark	MG	Madagascar	us	United States of America
Spain	MI.	Mali	VN	Vict Nam
Finland	MN	Mongolia		
	Australia Barhados Belgium Burkina Faso Bulgaria Benin Brazil Canada Central African Republic Congo Switzerland Côte d'Ivoire Cameroon Czechoslovakia Czech Republic Germany Denmark Spain	Australia GA Barhados GB Belgium GN Burkina Faso GR Bulgaria HU Benin IE Brazil IT Canada JP Central African Republic KP Congo Switzerland KR Côte d'Ivoire KZ Canteroon Li Czechoslovakia LK Czech Republic LU Germany MC Denmark MG Spain MI.	Australia GA Gabon Barhados GB United Kingdom Belgium GN Guinea Burkina Faso GR Greece Bulgarla HU Hungary Benin IE Ireland Brazil IT Italy Canada JP Japan Central African Republic KP Democratic People's Republic Congo of Korea Switzerland KR Republic of Korea Côte d'Ivoire KZ Kazakhstan Cameroon LI Licchtenstein Czechoslovakia LK Sri Lanka Cyech Republic LU Luxembourg Germany MC Monaco Denmark MG Madagascar Spain MI. Mali	Australia GA Gabon MW Barhados GB United Kingdom NL Belgium GN Guinea NO Burkina Faso GR Greece NZ Bulgarla HU Hungary PL Benin IE Ireland PT Brazil IT Italy RO Canada JP Japan RU Central African Republic KP Democratic People's Republic SD Congo of Korea SE Switzerland KR Republic of Korea SK Côte d'Ivoire KZ Kazakhstan SN Cameroon LI Licehtenstein SU Czechoslovakia LK Sri Lanka TD Czech Republic LU Lucembourg TG Germany MC Monaco UA Denmark MG Madagascar US Spain MI Mali Mali

WO 93/25517 PCT/GB93/01266

1

Trisubstituted phenyl derivatives as selective phosphodiesterase IV inhibitors

FIELD OF THE INVENTION

10

20

25

This invention relates to a novel series of trisubstituted phenyl derivatives, to processes for their preparation, to pharmaceutical compositions containing them, and to their use in medicine.

BACKGROUND TO THE INVENTION

Many hormones and neurotransmitters modulate tissue function by elevating intra-cellular levels of adenosine 3', 5'-cyclic monophosphate (cAMP). The cellular levels of cAMP are regulated by mechanisms which control synthesis and breakdown. The synthesis of cAMP is controlled by adenylyl cyclase which may be directly activated by agents such as forskolin or indirectly activated by the binding of specific agonists to cell surface receptors which are coupled to adenylyl cyclase. The breakdown of cAMP is controlled by a family of phosphodiesterase (PDE) isoenzymes, which also control the breakdown of guanosine 3',5'-cyclic monophosphate (cGMP). To date, seven members of the family have been described (PDE I-VII) the distribution of which varies from tissue to tissue. This suggests that specific inhibitors of PDE isoenzymes could achieve differential elevation of cAMP in different tissues, [for reviews of PDE distribution, structure, function and regulation, see Beavo & Reifsnyder (1990) TIPS, 11: 150-155 and Nicholson et al (1991) TIPS, 12: 19-27].

There is clear evidence that elevation of cAMP in inflammatory leukocytes leads to inhibition of their activation. Furthermore, elevation of cAMP in airway smooth muscle has a spasmolytic effect. In these tissues, PDE IV plays a major role in the hydrolysis of cAMP. It can be expected, therefore, that selective inhibitors of PDE IV would have therapeutic effects in inflammatory diseases such as asthma, by achieving both anti-inflammatory and bronchodilator effects.

The design of PDE IV inhibitors has met with limited success to date, in that many of the potential PDE IV inhibitors which have been synthesised have lacked potency and/or have been capable of inhibiting more than one type of PDE isoenzyme in a non-selective manner. Lack of a selective action has been a particular problem given the widespread role of cAMP in vivo and what is needed are potent selective PDE IV inhibitors with an inhibitory action against PDE IV and little or no action against other PDE isoenzymes.

SUMMARY OF THE INVENTION

10

15

20

We have now found a novel series of trisubstituted phenyl derivatives, members of which are potent inhibitors of PDE IV at concentrations at which they have little or no inhibitory action on other PDE isoenzymes. These compounds inhibit the isolated PDE IV enzyme and also elevate cAMP in isolated leukocytes. Certain compounds prevent inflammation in the lungs induced by carrageenan, platelet-activating factor (PAF), interleukin-5 (IL-5) or antigen challenge. These compounds also suppress the hyperresponsiveness of airway smooth muscle seen in inflamed lungs. The compounds of the invention are therefore of use in medicine, especially in the prophylaxis and treatment of asthma.

Thus according to one aspect of the invention, we provide a compound of formula (1)

$$X \longrightarrow \mathbb{R}^2$$

$$Z(CH_2)_n \mathbb{R}^4$$
(1)

wherein

Y is a halogen atom or a group -OR1 where R1 is an optionally substituted alkyl group;

WO 93/25517 3 PCT/GB93/01266

R² is an optionally substituted cycloalkyl, cycloalkenyl or polycycloalkyl group;

Z is a group $-N(R^3)CO$ - or $-CON(R^3)$ - where R^3 is a hydrogen atom or an alkyl, anyl or aralkyl group;

5 R4 is an aryl or heteroaryl group;

25

X is -O-, -S- -CH₂- or -N(\mathbb{R}^5)-, where \mathbb{R}^5 is a hydrogen atom or an alkyl group:

n is zero or an integer of value 1, 2 or 3; and the salts, solvates and hydrates thereof.

10 In the compounds of formula (1), when Y is a halogen atom it may be for example a fluorine, chlorine, bromine or iodine atom.

When Y in the compounds of formula (1) is a group OR¹, R¹ may be, for example, an optionally substituted straight or branched alkyl group, for example, a C₁₋₆ alkyl group, e.g. a C₁₋₃ alkyl group, such as a methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl, t-butyl, n-pentyl, i-pentyl or n-hexyl group. Optional substituents which may be present on R¹ groups include one or more halogen atoms, e.g. fluorine, chlorine, bromine or iodine atoms.

When R^2 in the compounds of formula (1) is an optionally substituted cycloalkyl or cycloalkenyl group it may be for example a C_{3-8} cycloalkyl group such as a cyclobutyl, cyclopentyl or cyclohexyl group or a C_{3-8} cycloalkenyl group such as a cyclobutenyl, cyclopentenyl or cyclohexenyl group, each cycloalkyl or cycloalkenyl group being optionally substituted by one, two or three substituents selected from halogen atoms, e.g. fluorine, chlorine, bromine or iodine atoms, C_{1-6} alkyl e.g. C_{1-3} alkyl such as methyl or ethyl, hydroxyl or C_{1-6} alkoxy e.g. C_{1-3} alkoxy such as methoxy or ethoxy groups. Polycycloalkyl groups represented by the group R^2 include optionally substituted C_{7-10} polycycloalkyl groups e.g. C_{7-10} bicycloalkyl or C_{7-10} tricycloalkyl groups such as bicyclo[2.2.1] heptyl or indanyl groups optionally substituted by one, two or three substituents as just described for substituted cycloalkyl or cycloalkenyl groups represented by R^2 .

The group R^3 in the compounds of formula (1) may be a hydrogen atom or a straight or branched C_{1-6} alkyl group, e.g. a C_{1-3} alkyl group. Particular groups include methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl, or t-

4 WO 93/25517 PCT/GB93/01266

butyl groups. Alternatively, R3 may represent a C₆₋₁₂ aryl group such as an optionally substituted phenyl group or a C₆₋₁₂ aryl C₁₋₆ alkyl group, e.g. an optionally substituted phenylC1-3alkyl group such as an optionally substituted benzyl or phenethyl group.

5 When the group R4 in compounds of formula (1) is an aryl group, it may be an optionally substituted C₆₋₁₂ aryl group such as an optionally substituted phenyl or 1- or 2-naphthyl group. Heteroaryl groups represented by R4 include optionally substituted C₃₋₆ heteroaryl groups containing for example one or two heteroatoms selected from -O- or -S- or groups -N(R5)- [where R5 is as defined above], such as pyrrolyl, furanyl, oxazolyl, thiazolyl, pyrazoyl, pyridinyl or pyrimidinyl groups. The heteroaryl group may be attached to the remainder of the molecule of formula (1) through any ring carbon or heteroatom as desired.

The aryl or heteroaryl groups represented by R3 and R4 in compounds of formula (1) may each optionally be substituted by one, two or more substituents (R6) selected from halogen atoms, e.g. fluorine, chlorine, bromine or iodine atoms, or C₁₋₆ alkyl, e.g. methyl or ethyl, C₁₋₆ alkoxy, e.g. methoxy or ethoxy, C₂₋₆ alkylenedioxy, e.g. ethylenedioxy, C₅₋₇ cycloalkoxy, e.g. cyclopentyoxy, haloC₁₋₆ alkyl, e.g. tri-fluoromethyl, C₁₋₆ alkylamino, e.g. methylamino or ethylamino, C₁₋₆ dialkylamino, e.g. dimethylamino or diethylamino, amino (NH₂), nitro, cyano, hydroxyl (OH), carboxyl (CO₂H), -CO₂R⁷ [where R⁷ is a C₁₋₆ alkyl e.g. methyl or ethyl, C₆₋₁₂ arylC₁₋₃ alkyl, e.g. benzyl or phenethyl or C₆₋₁₂ aryl, e.g. phenyl group], C₁₋₆ alkanoyl e.g. acetyl, sulphonyl (-SO₃H), C₁₋₆ alkylsulphonyl, e.g. methylsulphonyl, aminosulphonyl (-SO₂NH₂), C₁₋₆ alkylaminosulphonyl e.g. methylaminosulphonyl or ethylaminosulphonyl, C₁₋₆ dialkylaminosulphonyl, e.g. dimethylaminosulphonyl or diethylaminosulphonyl, carboxamido (-C₁₋₆ alkylaminocarbonyl, e.g. methylaminocarbonyl or CONH₂), ethylaminocarbonyl, C₁₋₆ dialkylaminocarbonyl, e.g. dimethylaminocarbonyl diethylaminocarbonyl, sulphonylamino (-NHSO₂H), alkylsulphonylamino, e.g. methylsulphonylamino or ethylsulphonylamino, C₁₋₆ dialkylsulphonylamino, e.g. dimethylsulphonylamino or diethylsulphonylamino groups, -NHCOR8 (where R8 is a C₁₋₆ alkyl group, e.g. a methyl, ethyl, n-propyl, or i-propyl group, or an aryl, e.g. phenyl or aryl 35 C₁₋₃ alkyl, e.g. benzyl, or phenethyl group), -NHCSR⁸, -NHCONH₂,

20

25

30

WO 93/25517 PCT/GB93/01266

5

-NHCONH, -NHCONHR⁷, -NHCON(R⁷)₂₄ (where each R⁷ group is the same or different), or C_{1-6} alkanoylamino C_{1-6} alkyl, e.g. acetylaminomethyl groups.

It will be appreciated that where two or more R⁶ substitutents are present, these need not necessarily be the same atom and/or groups. The R⁶ substituents may be present at any ring carbon atom away from the atom attached to the rest of the molecule of formula (1). Thus for example, in substituted phenyl groups represented by R³ or R⁴ any substituent(s) may be present at the 2-; 3-; 4-; 5-; 6-; 2,6-; 2,3- or 2,4- positions relative to the ring carbon atom attached to the remainder of the molecule. In another example, when the group R⁴ is a heteroaryl group any carbon atom not attached to the remainder of the molecule may be substituted. One, two or more carbon atoms may have substituents. Thus, in one example, when R⁴ is a substituted 4-pyridinyl group any substituent(s) may be present at the 2-; 3-; 5-; 6-; 2,5- or 3,5 positions relative to the nitrogen atom.

When the group -N(R^5)- is present in the compounds of formula (1), R^5 may be a hydrogen atom or a C_{1-6} alkyl group such as a methyl or ethyl group.

The presence of certain substituents in the compounds of formula (1) may enable salts of the compounds to be formed. Suitable salts include pharmaceutically acceptable salts, for example acid addition salts derived from inorganic or organic acids, and salts derived from inorganic and organic bases.

20

25

30

Acid addition salts include hydrochlorides, hydrobromides, hydroiodides, p-toluenesulphonates, phosphates, sulphates, acetates, trifluoroacetates, propionates, citrates, maleates, fumarates, malonates, succinates, lactates, oxalates, tartrates and benzoates.

Salts derived from inorganic or organic bases include alkali metal salts such as sodium or potassium salts, alkaline earth metal salts such as magnesium or calcium salts, and organic amine salts such as morpholine, piperidine, dimethylamine or diethylamine salts.

One particular group of compounds according to the invention has the formula (1) wherein Z is a -N(R³)CO- group and Y, R², R³, R⁴, X and n are as defined for formula (1), and the salts, solvates and hydrates thereof. Particular compounds of this type and those wherein Y is an -OR¹ group, R² is an optionally substituted cycloalkyl or polycycloalkyl group and R¹, R³, R⁴, X and n are as defined for formula (1) and the salts, solvates and hydrates thereof.

Another group of compounds according to the invention has the formula (1) wherein Z is a -CON(R³)- group and Y, R², R³, R⁴, X and n are as defined for formula (1), and the salts, solvates and hydrates thereof. Particular compounds of this type are those wherein Y is an -OR¹ group, R² is an optionally substituted cycloalkyl or polycycloalkyl group and R¹, R³, R⁴ X and n are as defined for formula (1) and the salts, solvates and hydrates thereof.

The following preferences apply alternatively to the compounds of formula (1) when Z is either a -N(R³)CO- or a -CON(R³)- group.

In the compounds of formula (1), the group Y is preferably an -OR 1 group. Particularly useful compounds of this type are those where R 1 is an optionally substituted C $_{1-3}$ alkyl group, particularly an ethyl group, or especially a methyl group.

The group X in compounds of formula (1) may be for example -S-, -CH₂- or -N(\mathbb{R}^5)-, but is preferably -O-.

R² in compounds of formula (1) is preferably an optionally substituted cyclopentyl group.

25 In one group of compounds of formula (1) n is preferably zero.

20

30

In another group of compounds of formula (1) n is an integer of value 1, 2 or 3, and is preferably an integer of value 1.

A useful group of compounds according to the invention has the formula (1) wherein Y is a group $-OR^1$ where R^1 is an optionally substituted straight or branched C_{1-3} alkyl group, R^2 is an optionally substituted C_{3-8} cycloalkyl

group, Z is either -N(R³)CO- or -CON(R³)- where R³ is a hydrogen atom, R⁴ is a C_{6-12} aryl or C_{3-6} heteroaryl group, X is -O-, n is zero or an integer of value 1 and the salts, solvates and hydrates thereof. In compounds of this type, n is, in particular, zero.

5 A particularly useful group of compounds of formula (1) has the formula (2):

$$CCH_3$$
 $CCH_2)_nR^4$
(2)

and the salts, solvates and hydrates thereof, wherein Z, n, and R⁴ are as defined for formula (1).

Particular compounds of formula (2) are those wherein n is an integer of value 1, or, especially, is zero.

In the compounds of formulae (1) and (2), Z is preferably - NHCO- or -CONH-.

R4 in the compounds of formulae (1) and (2) is preferably an optionally substituted phenyl, or an optionally substituted 2-,3- or, especially, 4-pyridinyl group. Particularly useful substituents include halogen atoms such as fluorine, chlorine, or bromine atoms, and groups selected from nitro, trifluoromethyl, carboxamido, -CO₂CH₃, -CH₃, or -OCH₃.

Particularly useful compounds according to the invention are:

3-Cyclopentyloxy-4-methyloxy-N-(2-Nitrobenzoyl)aniline;

N-(3-Cyclopentyloxy-4-methyloxyphenyl)-4-pyridinecarboxamide;

20 3-Cyclopentyloxy-4-methyloxy-N-(3-Nitrobenzoyl)aniline;

N-(3-Cyclopentyloxy-4-methyloxyphenyl)-1,2-benzenedicarboxamide;

N-Benzoyl-3-cyclopentyloxy-4-methyloxyaniline;

15

WO 93/25517 8 PCT/GB93/01266

Methyl 2-[N-(3-Cyclopentyloxy-4-methyloxyphenyl)carbamoyl]benzoate; N-(3-Cyclopentyloxy-4-methyloxyphenyl)-3,5-dichloro-4-

pyridinecarboxamide;

20

N-Phenyl-3-cyclopentyloxy-4-methyloxybenzamide;

5 N-(2-Nitrophenyl)-3-cyclopentyloxy-4-methoyloxybenzamide;

N-(3,5-Dichloropyrid-4-yl)-3-cyclopentyloxy-4-methyloxybenzamide;

N-(2,6-Difluorophenyl)-3-cyclopentyloxy-4-methyloxybenzamide;

N-Pyrimidin-4-yl-3-cyclopentyloxy-4-methoxybenzamide;

N-Pyridin-4-yl-3-cyclopentyloxy-4-methyloxybenzamide;

10 N-Pyrizin-2-yl-3-cyclopentyloxy-4-methyloxybenzamide;

N-Benzyl-3-cyclopentyloxy-4-methyloxybenzamide; and the salts, solvates and hydrates thereof.

Compounds according to the invention are selective and potent inhibitors of PDE IV. The ability of the compounds to act in this way may be simply determined by the tests described in the Examples hereinafter.

The compounds according to the invention are thus of particular use in the prophylaxis and treatment of human diseases where an unwanted inflammatory response or muscular spasm is present and where the elevation of cAMP levels may be expected to prevent or alleviate the inflammation and relax the muscle.

Particular uses to which the compounds of the invention may be put include the prophylaxis and treatment of asthma, especially inflamed lung associated with asthma, or in the treatment of inflammatory airway disease, chronic bronchitis, eosinophilic granuloma, psoriasis and other benign and malignant proliferative skin diseases, endotoxic shock, septic shock, ulcerative colitis, Crohn's disease, reperfusion injury of the myocardium and brain, chronic glomerulonephritis, atopic dermatitis, urticaria, allergic rhinitis, adult respiratory distress syndrome, diabetes insipidus, allergic conjunctivitis and vernal conjunctivitis.

For the prophylaxis or treatment of disease the compounds according to the invention may be administered as pharmaceutical compositions, and according to a further aspect of the invention we provide a pharmaceutical composition which comprises a compound of formula (1) together with one or more pharmaceutically acceptable carriers, excipients or diluents.

WO 93/25517 PCT/GB93/01266

Compounds according to the invention may also elevate cAMP in lymphocytes and thereby suppress unwanted lymphocyte activation in immune-based diseases such as rheumatoid arthritis, transplant rejection and graft versus host disease.

Pharmaceutical compositions according to the invention may take a form suitable for oral, buccal, parenteral or nasal administration, or a form suitable for administration by inhalation or insufflation.

10

15

20

25

30

For oral administration, the pharmaceutical compositions may take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g. pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g. lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (e.g. magnesium stearate, talc or silica); disintegrants (e.g. potato starch or sodium glycollate); or wetting agents (e.g. sodium lauryl sulphate). The tablets may be coated by methods well known in the art. Liquid preparations for oral administration may take the form of, for example, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents, emulsifying agents, non-aqueous vehicles and preservatives. The preparations may also contain buffer salts, flavouring, colouring and sweetening agents as appropriate.

Preparations for oral administration may be suitably formulated to give controlled release of the active compound.

For buccal administration the compositions may take the form of tablets or lozenges formulated in conventional manner.

The compounds of formula (1) may be formulated for parenteral administration by injection e.g. by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form. The compositions for injection may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents

WO 93/25517 10 PCT/GB93/01266

such as suspending, stabilising and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g. sterile pyrogen-free water, before use.

In addition to the formulations described above, the compounds of formula (1) may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation or by intramuscular injection.

For nasal administration or administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation for pressurised packs or a nebuliser, with the use of suitable propellant, e.g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas.

The compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the active ingredient. The pack or dispense device may be accompanied by instructions for administration.

The quantity of a compound of the invention required for the prophylaxis or treatment of a particular inflammatory condition will vary depending on the compound chosen, and the condition of the patient to be treated. In general, however, daily dosages may range from around 0.01mg/kg to 100mg/kg, e.g. around 1mg/kg to 40mg/kg body weight for oral or buccal administration from around 0.001mg/kg to 50mg/kg body weight for parenteral administration, and around 5mg to around 1000mg for nasal administration or administration by inhalation or insufflation.

20

The compounds according to the invention may be prepared by the following processes. The symbols Y, R², R³, R⁴ and X when used in the formulae below are to be understood to represent those groups described above in relation to formula (1) unless otherwise indicated.

Thus according to another aspect of the invention, a compound of formula (1) may be prepared either (a) by coupling an amine of formula (3a):

$$X - R^2$$

$$R^3 - NH$$
(3a)

with an acid $R^4(CH_2)_nCO_2H$ or an active derivative thereof or (b) by coupling an acid of formula (3b)

$$X - R^2$$
 CO_2H
(3b)

or an active derivative thereof with an amine $R^4(CH_2)_nNHR^3$. Active derivatives of acids of formula (3b) or acids $R^4(CH_2)_nCO_2H$ include, for example, acid anhydrides, or acid halides, such as acid chlorides.

The coupling reactions may be performed using standard conditions for reactions of this type. Thus for example, the reaction may be carried out in a solvent, for example an inert organic solvent such as an ether, e.g. a cyclic ether such as tetrahydrofuran, an amide, e.g. a substituted amide such as dimethylformamide, or a halogenated hydrocarbon such as dichloromethane, at a low temperature, e.g. -30°C to ambient temperature such as -20°C to 0°C, optionally in the presence of a base, e.g. an organic base such as an amine, e.g. triethylamine or a cyclic amine such as N-methylmorpholine. Where an acid of formula (3b) or R4(CH2)CO2H is used, the reaction may additionally be performed in the presence of a condensing agent, for example a diimide such as N, N'-dicyclohexylcarbodiimide, advantageously in the presence of a triazole such as 1-hydroxybenzotriazole. Alternatively, the acid may be reacted with a

10

15

chloroformate, for example ethylchloroformate, prior to reaction with the amine of formula (3).

Intermediate amines of formula (3a) in which R³ is a hydrogen atom may be prepared by hydrogenation of a corresponding nitro compound of formula (4):

$$X \longrightarrow \mathbb{R}^2$$
 NO_2
(4)

using for example hydrogen and a metal such as palladium or platinum optionally on a support such as carbon, in a solvent such as an alcohol e.g. methanol or ethanol, preferably at an elevated pressure and temperature

Intermediates of formula (3a) wherein R³ is as defined above other than a hydrogen atom may be prepared by reaction of a compound of formula (3a) where R³ is a hydrogen atom with an appropriate halide R³Hal [where Hal is a halogen atom such as a chlorine, bromine or iodine atom] in the presence of a base such as sodium hydride or an alkoxide such as sodium ethoxide, in a solvent such as an alcohol e.g. ethanol, at ambient temperature or above e.g. around 40°C or 50°C.

Intermediates of formula (4) in which X is -O-, -S- or -N(\mathbb{R}^5)- may be prepared by alkylation of a corresponding compound of formula (5):

15

using a reagent R²Hal [where Hal is as just defined]. The reaction conditions may be those just described for the alkylation of intermediates of formula (3a), or, alternatively, where for example the group X is -O- a base such as an alkali metal hydroxide or carbonate, e.g. cesium carbonate may be used in an inert solvent such dimethylformamide at ambient temperature.

Intermediates of formula (4) in which X is -CH₂-, may be prepared by reaction of a compound of formula (6)

with an acid halide R²COHal in the presence of a catalyst such as aluminium chloride followed by reduction using zinc and an inorganic acid such as hydrochloric acid, or hydrazine and a base such as an alkali metal hydroxide, e.g. sodium hydroxide.

Intermediates of formulae (5) and (6) and the reagents R⁴(CH₂)_nCO₂H and R⁴Hal are either known compounds or may be prepared from known starting materials by methods analogous to those used for the preparation of the known compounds.

Intermediate acids of formula (3b) may be prepared by hydrolysis of a corresponding ester of formula (7):

10

$$X - R^2$$

$$OR^9$$
(7)

where R⁹ is an alkyl group such as a methyl group, by heating in the presence of a base, for example an alkali metal hydroxide such as lithium hydroxide in a solvent such as an alcohol, e.g. methanol.

Intermediates of formula (7) in which X is -O-, -S- or -N(R⁵)- may be prepared by alkylation of a corresponding compound of formula (8):

using a reagent R²Hal [where Hal is as just defined] using the reagents and conditions as just described for the preparation of intermediates of formula (5).

Intermediates of formula (7) in which X is -CH₂-, may be prepared by reaction of a compound of formula (9):

with an acid halide R²COHal as described above for the preparation of compounds of formula (6)

Intermediates of formulae (8) or (9) may be prepared by esterification of the corresponding acids of formulae (10) or (11):

using for example acetyl chloride and an alcohol R⁹OH at an elevated temperature.

Intermediate acids of formulae (10) and (11) and the reagents $R^4(CH_2)_nNHR^3$, R^2Hal and R^9OH are either known compounds or may be prepared from known starting materials by methods analogous to those used for the preparation of the known compounds.

10

Salts of compounds of formula (1) may be prepared by reaction with an appropriate acid or base in a suitable solvent using conventional procedures.

DESCRIPTION OF SPECIFIC EMBODIMENTS

The following Examples illustrate the invention.

INTERMEDIATE 1

3-Cyclopentyloxy-4-methyloxy-1-nitrobenzene

5 2-Methyloxy-5-nitrophenol (14.27g, 84.4mmol) was dissolved in dry dimethylformamide (80ml). Cyclopentylbromide (176g, 127ml, 118.2mmol) and cesium carbonate (38.5g, 118.2mmol) were added and the suspension stirred overnight at room temperature. The mixture was poured into saturated Na₂CO₃ (200ml) and extracted with ethyl acetate (700ml). The organic layer was washed with saturated Na₂CO₃ until no yellow colour was detected. The organic layer was then dried (MgSO₄) and concentrated in vacuo to give the title compound as a yellow solid (15.0g).

¹H NMR δ (CDCl₃) 7.85 (1H, dd), 7.72 (1H, d), 6.90 (1H, d), 4.82-4.92 (1H, m), 3.92 (3H, s), 1.60-2.10 (8H, m).

15 INTERMEDIATE 2

20

3-Cyclopentyloxy-4-methyloxyaniline

Intermediate 1 was dissolved in degassed methanol and 10% Pd/C (450mg) was added. The mixture was stirred under H_2 gas at room temperature overnight. The catalyst was removed by filtration and the methanol concentrated in vacuo to give the <u>title compound</u> as a dark oil (4.3g).

¹H NMR δ (CDCl₃·) 6.70 (1H, d), 6.27 (1H, d), 6.20 (1H, dd), 4.68-4.78 (1H, m), 3.75 (3H, s), 3.42 (1H, br s), 1.50-1.95 (8H, m).

INTERMEDIATE 3

3-Hydroxy-4-methoxy-O-methylbenzoate

To methanol (150ml) at 0°C was added acetyl chloride (5.6g, 5.07ml). After 10 minutes a suspension of 3-Hydroxy-4-methoxybenzoic acid (10g) was added. The reaction mixture was refluxed for 6 hours, cooled and concentrated in vacuo. The residue was dissolved in H₂O, washed with NaHCO₃ (2x100ml), dried MgSO₄ and concentrated to give the title compound as a crystalline solid (9.8g).

¹H NMR δ (CDCl₃) 7.68 (1H, dd), 7.55(1H, d), 6.85 (1H, d), 4.80-4.91 (1H, m), 4.90 (3H, s), 4.88 (3H, s), 1.55-2.08 (8H, m).

INTERMEDIATE 4

15

25

3-Cyclopentyloxy-4-methyloxy-0-methylbenzoate

Intermediate 3 was refluxed in dry acetone (20ml) in the presence of K_2CO_3 (1.65g) and cyclopentylbromide (1.77g, 1.6ml) for 48 hours. The mixture was poured into ether/Na₂CO₃. The ether layer was removed and washed with Na₂CO₃ (3x100ml). The organic layer was dried over MgSO₄ and concentrated to give a white solid, recrystallisation of which from $H_2O/hexane$ gave the <u>title compound</u> (1.54g) as a white solid.

¹H NMR δ (CDCl₃) 7.62 (1H, dd), 7.57 (1H, d), 6.85 (1H, d), 4.80-4.90 (1H, m), 3.90 (3H, s), 3.88 (3H, s), 1.55-2.08 (8H, m).

INTERMEDIATE 5

3-Cyclopentyloxy-4-methyloxy benzoic acid

Intermediate 4 (1.18g) was added to CH₂Cl₂/methanol/H₂O (15:5:5, 25ml) and heated to reflux for 2 hours. The reaction mixture was cooled and poured into 1.0m HCl (50ml), then extracted with ethyl acetate (50ml). The

WO 93/25517 18 PCT/GB93/01266

organic layer was removed, washed with 1.0M HCI (3x50ml), dried (MgSO₄) and concentrated in vacuo to give the <u>title compound</u> (900mg) as a white solid which was recrystallised twice from ethyl acetate/hexane to give a white solid (450mg).

¹H NMR δ (CDCl₃): 7.75 (1H, dd), 7.60 (1H, d), 6.88 (1H, d), 4.70-4.80 (1H, m), 3.92 (3H, s), 1.55-2.05 (8H, m).

EXAMPLE 1

10

15

20

25

N-Benzoyl-3-cyclopentyloxy-4-methyloxyaniline

To an ice cooled solution of Intermediate 2 (20g, 9.66mmol) in anhydrous pyridine (20ml) was added benzoyl chloride dropwise (2.04g, 1.68ml, 14.5mmol) followed by N,N-dimethylaminopyridine (5mg). The reaction mixture was stirred at room temperature for 3 hours, poured into 1.0M HCl (100ml) and extracted with ethyl acetate. The ethyl acetate layer was washed with 1.0M HCl (2x50ml) and dried over MgSO₄. Concentration in vacuo gave a brownish oil which was recrystallised twice from CH₂Cl₂/hexane to give the title compound as a white solid (1.0g).

¹H NMR δ (CDCl₃) 7.80-7.95 (3H, m), 7.40-7.55 (4H, m), 6.98 (1H, dd),6.82 (1H, d), 4.72-4.82 (1H, m), 3.83 (3H, s), 1.50-2.05 (8H, m).

The following compounds were prepared in a similar manner to the compound of Example 1 using Intermediate 2 and the approprirate chloride:

EXAMPLE 2

3-Cyclopentyloxy-4-methyloxy-N-(3-Nitrobenzoyl)aniline

¹H NMR δ (CDCl₃) 8.70 (1H, br s), 8.36 (1H, d), 8.28 (1H, d), 8.20 (1H, s), 7.68 (1H, t), 7.41 (1H, s), 7.07 (1H, dd), 6.85 (1H, d), 4.70-4.82 (1H, m), 3.85 (3H, s), 1.50-2.05 (8H, m).

WO 93/25517 PCT/GB93/01266

EXAMPLE 3

3-Cyclopentyloxy-4-methyloxy-N-(2-Nitrobenzoyi)aniline

¹H NMR δ (CDCl₃) 8.10 (1H, d), 7.60-7.80 (3H, m), 7.43 (1H, s), 7.35 (1H, d), 6.90 (1H, dd), 6.85 (1H, d), 4.75-4.85 (1H, m), 3.87 (3H, s), 1.50-2.05 (8H, m).

EXAMPLE 4

3-Cyclopentyloxy-4-methyloxy-N-(4-Nitrobenzoyl)aniline

¹H NMR δ (CDCl₃) 8.35 (2H, d), 8.02 (2H, d), 7.85 (1H, br s), 7.45 (1H, d), 7.05 (1H, dd), 6.87 (1H, d), 4.75-4.85 (1H, m), 3.87 (3H, s), 1.50-2.05 (8H, m).

EXAMPLE 5

10

15

20

25

N-(3-Cyclopentyloxy-4-methyloxyphenyl)-2-pyridinecarboxamide

¹H NMR δ (CDCl₃) 9.95 (1H, br s), 8.58 (1H, dd), 8.28 (1H, dd), 7.90 (1H, m), 7.62 (1H, d), 7.45 (1H, m), 7.12 (1H, dd), 6.88 (1H, d), 4.80-4.90 (1H, m), 3.87 (3H, s), 1.55-2.10 (8H, m).

EXAMPLE 6

N-(3-Cyclopentyloxy-4-methyloxyphenyl)-3-pyridinecarboxamide

¹H NMR δ (CDCl₃) 9.08 (1H, d), 8.78 (1H, dd), 8.20 (1H, dd), 8.05 (1H, br s), 7.35-7.45 (2H, m), 7.02 (1H, dd), 6.85 (1H, d), 4.75-4.85 (1H, m), 3.87 (3H, s), 1.55-2.10 (8H, m).

EXAMPLE 7

N-(3-Cyclopentyloxy-4-methyloxyphenyl)-4-pyridinecarboxamide

¹H NMR δ (CDCl₃) 8.76 (2H, d), 8.0 (1H, br s), 7.70 (2H, d), 7.45 (1H, d), 7.05 (1H, dd), 6.85 (1H, d), 4.75-4.85 (1H, m), 3.88 (3H, s), 1.55-2.05(8H, m).

EXAMPLE 8

N-(3-Cyclopentyloxy-4-methyloxyphenyl)-2-phenylacetamide
1H NMR δ (CDCl $_3$) 7.24-7.45 (6H, m), 7.02 (1H, br s), 6.68-6.80 (2H, m), 4.70-4.80 (1H, m), 3.80 (3H, s), 3.72 (2H, s), 1.55-2.05 (8H, m).

5 EXAMPLE 9

N-(3-Cyclopentyloxy-4-methyloxyphenyl)-1,2-benzenedicarboxamide 1H NMR δ (CDCl₃) 8.98 (H, s), 7.60 (1H, dd), 7.48 (1H, dd), 7.35-7.45 (3H, m), 7.05 (1H, dd), 6.78 (2H, br s + d), 5.90 (1H, br s), 4.70-4.80 (1H, m), 3.83 (3H, s), 1.50-2.05 (8H, m).

10 EXAMPLE 10

Methyl-2-[N-(3-Cyclopentyloxy-4-methyloxyphenyl)carbamoyi]benzoate 1 H NMR $_{0}$ (CDCl3) 7.92 (H, d), 7.48-7.65 (4H, m), 7.49 (1H, d), 7.0 (1H, dd), 6.83 (1H, d), 4.75-4.87 (1H, m), 3.88 (3H, s), 3.83 (3H, s), 1.5-2.0 (8H, m).

15 EXAMPLE 11

EXAMPLE 12

N-(3-Cyclopentyloxy-4-methyloxyphenyl)-N-methyl-2-nitrobenzamide

1H NMR δ (CDCl₃) 8.10 (1H, d), 7.55-7.75 (3H, m), 7.35 (1H, d), 6.98(1H, dd), 6.83 (1H, d), 4.78-4.85 (1H, m), 3.86 (3H, s), 2.7 (3H, s), 1.50-2.05 (8H, m).

WO 93/25517 PCT/GB93/01266

EXAMPLE 13

N-(3-Cyclopentyloxy-4-methyloxyphenyl)-3-cyclopentyloxy-4-methyloxybenzamide

¹H NMR δ (CDCl₃) 7.75 (1H, br s), 7.50 (2H, s), 7.38 (1H, dd), 6.8-7.0 (3H, m), 4.75-4.90 (2H, m), 3.90 (3H, s), 3.85 (3H, s), 1.50-2.05 (16H, m).

EXAMPLE 14

N-(3-Cyclopentyloxy-4-methyloxyphenyl)-N-phenyl-4-pyridinecarboxamide

¹H NMR δ (CDCl₃) 8.50 (2H, d, J = 6Hz), 7.27 (2H, d, J = 6Hz), 7.19-7.37 (5H, m), 6.60-6.75 (3H, m), 4.58 (1H, br s), 3.80 (3H, s), 1.62-1.85 (6H, m), 1.50-1.62 (2H, m).

EXAMPLE 15

15

20

N-Phenyl-3-cyclopentyloxy-4-methyloxybenzamide

Intermediate 5 (448mg) was dissolved in dry tetrahydrofuran (20ml). Nemethylmorpholine (191mg, 209µl) was added and the reaction left 30 minutes to activate. Neat aniline (510mg, 500µl) was added and the reaction stirred at room temperature overnight. The mixture was then poured into 1.0M HCl and ethyl acetate (50ml) was added. The organic layer was separateed, washed with 1.0M HCl (3x50ml), aqueous Na₂CO₃ (3x50ml) and dried (MgSO). Recrystallisation from ethyl acetate/hexane gave the title compound as a white solid (100g).

¹H NMR δ (CDCl₃) 7.77 (1H, br s), 7.64 (2H, d), 7.49 (1H, d), 7.30-7.40 (3H, m), 7.14 (1H, t), 6.88 (1H, d), 4.82-4.92 (1H, m), 3.90 (3H, s), 1.55-2.05 (8H, m).

The following compounds were prepared in a similar method to that described for the preparation of the compound of Example 15, from Intermediate 5 and the appropriate amine:

WO 93/25517 22 PCT/GB93/01266

EXAMPLE 16

N-(2-Nitrophenyl)-3-cyclopentyloxy-4-methyloxybenzamide

¹H NMR δ (CDCl₃) 11.34 (1H, br s), 9.01 (1H, d), 8.30 (2H, d), 7.72 (1H, t), 7.55-7.60 (2H, m), 7.22 (2H, t), 7.0 (1H, d), 4.85-4.95 (1H, m), 3.93 (3H, s), 1.55-2.10 (8H, m).

EXAMPLE 17

N-Benzyl-3-cyclopentyloxy-4-methyloxybenzamide

¹H NMR δ (CDCl₃) 7.42 (1H, d), 7.20-7.40 (6H, m), 6.85 (1H, d), 6.40 (1H, br s), 4.80-4.92 (1H, m), 4.65 (2H, d), 3.88 (3H, s), 1.48-2.05 (8H, m).

10 **EXAMPLE 18**

N-(3,5-dichloropyrid-4-yl)-3-cyclopentyloxy-4-methyloxybenzamide
1H NMR δ (CDCl₃) 8.55 (2H, s), 7.71 (1H, s), 7.48-7.52 (2H, m), 6.93 (1H, d, J = 8Hz), 4.85-4.95 (1H, m), 3.93 (3H, s), 1.60-2.10 (8H, br m).

EXAMPLE 19

N-(2,6-Difluorophenyl)-3-cyclopentyloxy-4-methyloxybenzamide

1H NMR δ (CDCl₃) 6.83-7.60 (7H, m), 4.84 (1H, m), 3.90 (1H, s), 1.55-2.07 (8H, m).

EXAMPLE 20

N-Pyrimidin-4-yl-3-cyclopentyloxy-4-methoxybenzamide

¹H NMR δ (CDCl₃) 8.87 (1H, s), 8.67 (1H, d d), 8.55 (1H, br s), 8.30-8.35 (1H, dd), 7.52 (1H, d), 7.4 (1H, dd), (2H, d + dd), 6.9-6.95 (1H, d), 4.82-4.85 (1H, m), 3.91 (3H, s), 1.55-2.0 (8H, m).

WO 93/25517 23 PCT/GB93/01266

EXAMPLE 21

N-Pyridin-4-yl-3-cyclopentyloxy-4-methyloxybenzamide

¹H NMR δ (CDCl₃) 8.5-8.7 (2H, dd), 8.02-8.15 (1H, br s), 7.59-7.60 (2H, dd), 7.39-7.40 (1H, dd), 7.47 (1H, s), 6.85-6.91 (1H, d), 4.3-4.4 (1H, m), 3.9 (3H, s), 1.55-2.05 (8H, m).

EXAMPLE 22

10

15

25

N-Pyrizin-2-yl-3-cyclopentyloxy-4-methyloxybenzamide

¹H NMR δ (CDCl₃) 9.7 (1H, d), 8.45-8.52 (1H, br s), 8.49 (1H, d), 8.25 (1H, dd), 7.49 (1H, d), 7.4 (1H, dd), 6.9-7.0 (1H, d), 4.85-4.95 (1H, m), 3.95 (3H, s), 0.8-2.2 (8H, m).

The activity and selectivity of compounds according to the invention was demonstrated in the following tests.

1. <u>Isolated Enzyme</u>

The potency and selectivity of the compounds of the invention was determined using a battery of distinct PDE isoenzymes as follows:

- i. PDE I, rabbit heart
- ii. PDE II, rabbit heart
- iii. PDE III, rabbit heart
- iv. PDE IV, HL60 cells.
- The enzymes were purified to kinetic homogeneity using standard chromatographic techniques.

Phosphodiesterase activity was assayed as follows. The reaction was conducted in 150 μ l of standard mixture containing (final concentrations): 50mM TES-NaOH buffer (pH 7.5), 10mM MgCl₂, 0.1 μ M [³H]-cAMP and vehicle or various concentrations of the test compounds. The reaction was initiated by addition of enzyme and

WO 93/25517 24 PCT/GB93/01266

conducted at 30°C for between 5 to 30 mins. The reaction was terminated by addition of 50µl 2% trifluoroacetic acid containing [14C]-5'AMP for determining recovery of the product. An aliquot of the sample was then applied to a column of neutral alumina and the [3H]-cAMP eluted with 10ml 0.1 TES-NaOH buffer (pH8). The [3H]-5'-AMP product was eluted with 2ml 2M NaOH into a scintillation vial containing 10ml of scintillation cocktail. Recovery of [3H]-5'AMP was determined using the [14C]-5'AMP and all assays were conducted in the linear range of the reaction.

Compounds according to the invention were able to inhibit the action of the PDE IV HL60 enzyme at concentrations at which they had little or no effect on the action of each of the other PDE isoenzymes. Thus, compounds of the Examples have approximate Ki values (Ki PDE IV HL60 at (1μM) in the pM-μM range, for example the compounds of Examples 1 and 3 have approximate Ki values of 391nM and 179nM respectively.

2. The Elevation of cAMP in Leukocytes

5

20

25

The effect of compounds of the invention on intracellular cAMP was investigated using human neutrophils or guinea pig eosinophils. Human neutrophils were separated from peripheral blood, incubated with dihydrocytochalasin B and the test compound for 10 min and then stimulated with FMLP. Guinea pig eosinophils were harvested by peritoneal lavage of animals previously treated with intra-peritoneal injections of human serum. Eosinophils were separated from the peritoneal exudate and incubated with isoprenaline and test compound. With both cell types, suspensions were centrifuged at the end of the incubation, the cell pellets were resuspended in buffer and boiled for 10 min prior to measurement of cAMP by specific radioimmunoassay (DuPont).

The most potent compounds according to the invention induced a concentration -dependent elevation of cAMP in neutrophils and/or eosinophils at concentrations in the range 1nM to 1μM.

5

25

30

3. Suppression of Leukocyte Function

Compounds of the invention were investigated for their effects on superoxide generation and chemotaxis of human neutrophils. Neutrophils were separated from peripheral blood, incubated with dihydrocytochalasin B for superoxide generation only and test compound prior to stimulation with FMLP. The most potent compounds caused a concentration-dependent inhibition of superoxide generation and chemotaxis at concentrations in the range 0.1nM to 1µM.

4. Relaxation of Constricted Airway Smooth Muscle in vitro

The effects of compounds of the invention on guinea-pig isolated tracheal smooth muscle were investigated. Isolated tracheal rings were suspended in organ baths and immersed in oxygenated Krebs' solution. The smooth muscle was contracted with sub-maximal concentrations of histamine or carbachol prior to the addition of increasing concentrations of test compound to the organ baths. The most potent compounds caused a concentration-dependent reversal of both histamine and carbachol-induced contractions at concentrations in the range 1nM to 100μM. The compounds were generally more potent in reversing histamine-induced tone than carbachol-induced tone.

20 5. Effects on Cardiac Muscle in vitro

Compounds of the invention have been tested for their effects on isolated cardiac muscle. Right atrial and papillary muscles were dissected out from the hearts of guinea pigs and suspended in organ baths for measuring the rate (chronotropic) of spontaneously beating atria and force (inotropic) of the electrically stimulated papillary muscle. In these preparations, selective PDE IV inhibitors such as rolipram do not have any direct effects whereas selective PDE III inhibitors such as milrinone have positive chronotropic and inotropic effects. The non-specific PDE inhibitor theophylline, which is used in asthma as a bronchodilator, also causes significant cardiovascular changes such as

tachycardia. Selective PDE IV inhibitors have advantage over theophylline, therefore, through reduced cardiovascular side effects. The most potent and selective compounds of the invention had no direct effects on the atrial and papillary muscles in vitro at concentrations up to 100µM but in combination with PDE III inhibitors, these inhibitors showed an enhancement of chronotropic and inotropic activity, typical of selective type IV inhibitors.

6. Anti-inflammatory Activity in vivo

5

10

15

20

25

30 -

Compounds of the invention have been tested in models of experimental pleurisy in the rat induced by carrageenan, platelet-activating factor (PAF), or interleukin-5 (IL-5). The compounds caused a dose-dependent reduction in carrageenan-induced accumulation of inflammatory exudate and total leukocyte numbers at 6h following oral (1-50mg/kg), intravenous (0.1-10mg/kg), or intraperitoneal (0.1-10mg/kg) administration. Compounds of the invention also reduced PAF or IL-5-induced pleural eosinophilia at 24h following oral (1-50mg/kg), intravenous (0.1-10mg/kg) or intraperitoneal (0.1-10mg/kg) dosing.

7. Anti-allergic Activity in vivo

Compounds of the invention have been tested for effects on an IgEmediated allergic pulmonary inflammation induced by inhalation of antigen by sensitised guinea pigs. Guinea pigs were initially sensitised ovalbumin under mild cyclophosphamide-induced immunosuppression, by intraperitoneal injection of antigen in combinations with aluminium hydroxide and pertussis vaccine. Booster doses of antigen were given two and four weeks later and at six weeks, animals were challenged with aerosolised ovalbumin whilst under cover of an intraperitoneally administered anti-histamine agent (mepyramine). After a further 48h, bronchial alveolar lavages (BAL) were performed and the numbers of eosinophils and other leukocytes in the BAL fluids were counted. The lungs were also removed for histological examination for inflammatory damage. Administration of compounds of WO 93/25517 27 PCT/GB93/01266

the invention (0.1-10mg/kg i.p.), up to three times during the 48h following antigen challenge, lead to a significant reduction in the eosinophilia and the accumulation of other inflammatory leukocytes. There was also less inflammatory damage in the lungs of animals treated with compounds of the invention.

8. Effects on Pulmonary Dynamics

5

10

15

Compounds of the invention have been tested for their effects on ozone-induced hyperreactivity of the airways of guinea pigs. Following the inhalation of ozone, guinea pigs become very much more sensitive to the bronchoconstrictor effects of inhaled histamine than naive animals. There is a pronounced shift to the left (10-30 fold) of the dose response curve to histamine and a highly significant increase in the maximum increase in pulmonary resistance. Compounds of the invention administered 1h prior to ozone by the intraperitoneal (0.01-1mg/kg) or oral (0.1-10mg/kg) route caused a dose-dependent inhibition of ozone-induced hyperreactivity.

In general, in our tests above, compounds of the invention have had no observed toxic effects when administered to animals at the doses shown.

CLAIMS

A compound of formula (1)

$$X - R^2$$

$$Z(CH_2)_n R^4$$
(1)

wherein

Y is a halogen atom or a group -OR1 where R1 is an optionally substituted alkyl group;

 $\ensuremath{\mathsf{R}}^2$ is an optionally substituted cycloalkyl, cycloalkenyl or polycycloalkyl group;

Z is a group -N(\mathbb{R}^3)CO- or -CON(\mathbb{R}^3)- where \mathbb{R}^3 is a hydrogen atom or an alkyl, anyl or analkyl group;

R4 is an aryl or heteroaryl group;

X is -O-, -S-, -CH2- or -N(\mathbb{R}^5)-, where \mathbb{R}^5 is a hydrogen atom or an alkyl group;

n is zero or an integer of value 1, 2 or 3; and the salts, solvates and hydrates thereof.

- 2. A compound according to Claim 1 wherein Z is a -N(R3)CO- group.
- 3. A compound according to Claim 2 wherein Z is a -NHCO- group.
- 4. A compound according to Claim 1 wherein Z is a -CON(R3)- group.
- 5. A compound according to Claim 4 wherein Z is a -CONH- group.
- 6. A compound according to any one of Claims 1 to 5 wherein Y is an -OR¹ group where R¹ is an optionally substituted alkyl group.

t

- 7. A compound according to Claim 6 where R¹ is an optionally substituted straight or branched C₁₋₃ alkyl group.
- 8. A compound according to Claim 7 where R¹ is a methyl group.
- 9. A compound according to any one of Claims 1 to 8 where n is zero or an integer of value 1.
- 10. A compound according to Claim 9 where n is zero.
- 11. A compound according to any one of Claims 1 to 10 where R² is an optionally substituted C₃₋₈ cycloalkyl group.
- 12. A compound according to Claim 11 where R² is a cyclopentyl group.
- 13. A compound according to any one of Claims 1 to 12 wherein R⁴ is an optionally substituted phenyl or pyridinyl group.
- 14. A compound according to any of the preceding claims wherein X is -O-.
- 15. A compound according to any one of Claims 1 to 13 wherein X is -S-, -CH₂- or -N(R⁵)- where R⁵ is a hydrogen atom or an alkyl group.
- 16. 3-Cyclopentyloxy-4-methyloxy-N-(2-Nitrobenzoyl)aniline; N-(3-Cyclopentyloxy-4-methyloxyphenyl)-4-pyridinecarboxamide; 3-Cyclopentyloxy-4-methyloxy-N-(3-Nitrobenzoyl)aniline; N-(3-Cyclopentyloxy-4-methyloxyphenyl)-1,2-benzenedicarboxamide; N-Benzoyl-3-cyclopentyloxy-4-methyloxyaniline; Methyl2-[N-(3-Cyclopentyloxy-4-methyloxyphenyl)carbamoyl] benzoate;
 - N-(3-Cyclopentyloxy-4-methyloxyphenyl)-3,5-dichloro-4-pyridine-carboxamide;
 - N-Phenyl-3-cyclopentyloxy-4-methyloxybenzamide:
 - N-(2-Nitrophenyl)-3-cyclopentyloxy-4-methoyloxybenzamide:
 - N-(3,5-Dichloropyrid-4-yl)-3-cyclopentyloxy-4-methyloxybenzamide:
 - N-(2,6-Difluorophenyl)-3-cyclopentyloxy-4-methyloxybenzamide:
 - N-Pyrimidin-4-yl-3-cyclopentyloxy-4-methoxybenzamide;

j

N-Pyridin-4-yl-3-cyclopentyloxy-4-methyloxybenzamide; N-Pyrizin-2-yl-3-cyclopentyloxy-4-methyloxybenzamide; N-Benzyl-3-cyclopentyloxy-4-methyloxybenzamide; and the salts, solvates, and hydrates thereof.

- 17. A pharmaceutical composition comprising a compound according to any one of Claims 1 to 16 and a pharmaceutically acceptable diluent, carrier or excipient.
- 18. A method for the prophylaxis or treatment of a human disease where an unwanted inflammatory response or muscular spasm is present and where the elevation of cAMP levels may be expected to prevent or alleviate the inflammation and relax the muscle, comprising the step of administering to a patient susceptible to, or having, said disease an amount of a pharmaceutical composition according to Claim 17 effective for elevating cAMP levels to prevent or alleviate said inflammation and relax said muscle.
- 19. A process for the preparation of a compound of formula (1)

$$X \longrightarrow R^2$$

$$Z(CH_2)_nR^4$$
(1)

wherein

Y is a halogen atom or a group -OR1 where R1 is an optionally substituted alkyl group;

R² is an optionally substituted cycloalkyl, cycloalkenyl or polycycloalkyl group;

Z is a group -N(R³)CO- or -CON(R³)- where R³ is a hydrogen atom or an alkyl, aryl or aralkyl group;

R4 is a n aryl or heteroaryl group;

X is -O-, -S-, -CH₂- or -N(\mathbb{R}^5)-, where \mathbb{R}^5 is a hydrogen atom or an alkyl group;

n is zero or an integer of value 1, 2 or 3; and the salts, solvates and hydrates thereof.

which comprises either (a) coupling an amine of formula (3a)-

$$X - R^2$$

$$R^3 - NH$$
(3a)

wherein

Y, R^2 , R^3 and X are as defined for formula (1), with an acid $R^4(CH_2)_nCO_2H$, where R^4 and n are as defined for formula (1) ,or an active derivative thereof or (b) by coupling an acid of formula (3b)

$$X \longrightarrow \mathbb{R}^2$$
 CO_2H (3b)

or an active derivative thereof with an amine R4(CH₂)_nNHR³.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 93/01266

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶								
		Classification (IPC) or to both National Cl	•					
Int.Cl	. 5 CO7C233/			07D213/82				
	C07C233/	29; C07C235/56;	C07D213/75; C0	07D239/42				
II. FIELDS	SEARCHED							
Minimum Documentation Searched?								
Classification System Classification Symbols								
Int.Cl	. 5	C07C ; C07D						
		Documentation Searched other to the Extent that such Documents a						
III. DOCU		D TO BE RELEVANT ⁹						
Category °	Citation of De	ocument, 11 with indication, where appropris	ate, of the relevant passages 12	Relevant to Claim No. ¹³				
A	WO,A,9 207 567 (SMITHKLINE BEECHAM CORP.) 14 May 1992							
	see page 26-32	·						
A	WO,A,9 31 Octol	1-18						
	see abstract, page 1, line 6 - line 10, example 1; page 11, table 1							
P,X	EP,A,O 6 5 Augus	1,4-14, 16-19						
	line 12	tract, page 2, line 19; examples, especially e 39, page 10, line 40,	page 9, line					
			•					
"A" document defining the general state of the art which is not considered to be of particular relevance "I later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the								
· "E" ear	dier document but publing date	invention "X" document of particular relevance; the cleannot be considered novel or cannot be						
"L" doc whi	limed invention							
	ution or other special re	rason (as specified) oral disclosure, use, exhibition or	cannot be considered to involve an inver- document is combined with one or more	tive step when the				
oth	er means		ments, such combination being obvious	to a person skilled				
	"P" document published prior to the international filing date but in the art. later than the priority date claimed "A" document member of the same patent family							
IV. CERTIFICATION								
Date of the	Actual Completion of	the International Search	Date of Mailing of this International Sec	urch Report				
	20 AUG	UST 1993	- 3. 09. 93					
Internations	il Searching Authority		Signature of Authorized Officer	······································				
	EUROPE	AN PATENT OFFICE	SEUFERT G.H.					

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.

GB 9301266 SA 74986

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on

The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

20/0 20/08/93

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
WO-A-9207567	14-05-92	AU-A-	9030691	26-05-92
WO-A-9116303	31-10-91	AU-A- EP-A-	7777491 0526535	11-11-91 10-02-93
EP-A-0497564	05-08-92	AU-A- WO-A-	1188192 9212961	27-08-92 06-08-92
				·
			٠	

2